




ENT Associates

Making sense of your family's hearing, sinus and throat problems

Neck Dissection Surgery

This is an operation to remove cancerous lymph nodes from one side of the neck. Traditionally a radical neck dissection was performed to remove the lymph nodes as well as other non-vital tissues from the neck, including salivary gland tissue, muscle, and veins. As the procedure has evolved, less radical forms of the surgery are usually performed, preserving uninvolved parts of the neck (modified radical, functional, or selective neck dissection). This leads to fewer long-term complications, better functionality, and shorter hospital stays. The surgery can be performed alone or in combination with other procedures to treat cancer of the head and neck.



Why do I need a neck dissection?


Chains of lymph nodes in the neck form a barrier to the spread of cancerous cells. By removing the lymph nodes on the affected side, possible sites of the spread of cancer are removed. Other tissues in the neck are also removed, as cancer can sometimes spread outside of the lymph nodes. Your surgeon will determine the extent of the operation that you will need.

How is the surgery performed?

Generally, the procedure is performed in a hospital setting under general anesthesia, and you will usually be admitted after surgery for a few days. The incision site will vary depending on the location of the group of lymph nodes to

be removed but some examples are shown below.

After the lymph nodes and surrounding tissues are removed, drains are usually placed in the neck to prevent build-up of fluid under the skin that occurs after surgery. This usually also improves wound healing. The doctor will determine how long the drains need to stay in place, but they are often removed after discharge.



What are the risks and possible complications of surgery?

The risks and potential complications will depend on the extent of the cancer and the extent of surgery that was required.

- ◆ Nerve injury: Certain nerves that are located in the neck may either be involved with the cancer or may be in close proximity to the tissues and lymph nodes that are being removed. This can result in weakness of the lower lip, tongue, and shoulder
- ◆ Bleeding: This may require a return to surgery to identify the source of bleeding and control it
- ◆ Chyle leak: this is leakage of a milky lymphatic fluid (**chyle**) from injury to a lymph channel that drains fat

(continued on back)

absorbed from the intestine into the blood vessels of your neck. It usually occurs only on the left side. It generally will heal itself on its own, and your doctor may advise a strict low fat diet until the chyle leak subsides

- ◆ Difficulty swallowing or speaking

Your doctor will also generally advise to avoid any heavy lifting or strenuous activity for several days after surgery. Other instructions will usually be provided to you upon discharge from the hospital.

Clearwater

Office: 727-441-3588

1330 South Fort Harrison • Clearwater, FL 33756

Countryside

Office: 727-791-1368

3190 McMullen Booth Rd • Clearwater, FL 33761

Dunedin

Office: 727-450-0560

646 Virginia Street, 3rd Floor • Dunedin, FL 34698

Facial Aesthetics Center

Office: 727-446-FACE (3223)

1320 South Fort Harrison • Clearwater, FL 33756

Largo

Office: 727-397-8551

8787 Bryan Dairy Road, Suite 170 • Largo, FL 33777

Port Richey

Office: 727-819-0368

11031 US 19, Suite 104 • Port Richey, FL 34668

St. Petersburg

Office: 727-767-0933

6540 4th Street North, Suite C • St. Petersburg, FL 33702

Tampa

Office: 813-925-5000

10810 Sheldon Road • Tampa, FL 33626

Trinity

Office: 727-247-1234

11320 State Road 54 • Trinity, FL 34655



ENT
Associates

Making sense of your family's
hearing, sinus and throat problems